

# $k$ –homogeneous latin trades

Richard Bean<sup>b</sup>, Hoda Bidkhori<sup>a</sup>, Maryam Khosravi<sup>c</sup>  
and  
E. S. Mahmoodian<sup>a</sup>

<sup>a</sup> Department of Mathematical Sciences  
Sharif University of Technology  
P. O. Box 11365–9415, Tehran, I. R. Iran

<sup>b</sup> Institute for Studies in Theoretical Physics and Mathematics,  
P. O. Box 19395-5746, Tehran, I. R. Iran

<sup>c</sup> Department of Mathematical Sciences  
Teacher’s Training University, Tehran, I. R. Iran

## Abstract

Let  $T$  be a partial latin square and  $L$  a latin square such that  $T \subset L$ . Then  $T$  is called a *latin trade*, if there exists a partial latin square  $T^*$  such that  $T^* \cap T = \phi$  and  $(L \setminus T) \cup T^*$  is a latin square. We call  $T^*$  a *disjoint mate* of  $T$ . A latin trade is called  *$k$ –homogeneous* if each row and each column contains exactly  $k$  elements, and each element appears exactly  $k$  times. The number of elements in a latin trade is referred to as its *volume*.

It is shown by Cavenagh, Donovan, and Drápal (2003 and 2004) that 3–homogeneous and 4–homogeneous latin trades of volume  $3m$  and  $4m$ , respectively, exist for all  $m \geq 3$  and  $m \geq 4$ , respectively. We show that  $k$ –homogeneous latin trades of volume  $km$  exist for all  $3 \leq k \leq 8$  and  $m \geq k$ . Also we show that for each given  $k \geq 3$  and for  $m \geq k$ , all  $k$ –homogeneous latin trades of volume  $km$  exist except possibly for finitely many  $m$ .

## 1 Introduction

A *latin square*  $L$  of order  $n$  is an  $n \times n$  array of the set, say  $N = \{0, 1, 2, \dots, n - 1\}$ , where each element of  $N$  appears exactly once in each row and exactly once in each

column. We can represent each latin square as a set of 3-tuples

$$L = \{(i, j; k) \mid \text{element } k \text{ is located in position } (i, j)\}.$$

A *transversal* of a latin square of order  $n$  is a set of  $n$  positions, no two in the same row or same column, containing each elements of the set  $N$  exactly once.

A *partial latin square*  $P$  of order  $n$  is an  $n \times n$  array of elements from the set  $N$ , where each element of  $N$  appears at most once in each row and at most once in each column. The set  $S_P = \{(i, j) \mid (i, j; k) \in P\}$  of the partial latin square  $P$  is called the *shape* of  $P$  and  $|S_P|$  is called the *volume* of  $P$ .

For each  $0 \leq r, c, e \leq n - 1$  we introduce the following sets:

$$C_P^c = \{k \mid (i, c; k) \in P\}, R_P^r = \{k \mid (r, j; k) \in P\} \text{ and } E_P^e = \{(i, j) \mid (i, j; e) \in P\}.$$

We call a partial latin square  $T$  of order  $n$  a *latin trade* if there exists a partial latin square  $T^*$  of order  $n$ , called a *mate* of  $T$ , such that:

$$S_T = S_{T^*}, \text{ and if } (i, j; k) \in T \text{ and } (i, j; k^*) \in T^*, \text{ then } k \neq k^*,$$

$$\text{for each } r, 0 \leq r \leq n - 1, \text{ we have } R_T^r = R_{T^*}^r; \text{ and}$$

$$\text{for each } c, 0 \leq c \leq n - 1, \text{ we have } C_T^c = C_{T^*}^c.$$

A latin trade is called *k-homogeneous* if:

$$\text{for each } r, 0 \leq r \leq n - 1, \text{ we have } |R_T^r| = 0 \text{ or } k;$$

$$\text{for each } c, 0 \leq c \leq n - 1, \text{ we have } |C_T^c| = 0 \text{ or } k; \text{ and}$$

$$\text{for each } e, 0 \leq e \leq n - 1, \text{ we have } |E_T^e| = 0 \text{ or } k.$$

A latin trade of volume 4 is called an *intercalate*. In Figure 1 an intercalate  $(T, T^*)$  is shown. The elements of  $T^*$  is written as subscripts in the same array as  $T$ .

0 <sub>1</sub>	1 <sub>0</sub>
1 <sub>0</sub>	0 <sub>1</sub>

Figure 1: An intercalate

For more background on latin trades see [2], [6], and [5], and concepts which are not defined here may be found in [1]. It is proved in [4] and [3], that 3-homogeneous Latin trades of volume  $3m$  exist for all  $m \geq 3$ , and 4-homogeneous latin trades of volume  $4m$  exist for all  $m \geq 4$ . We prove that for each given  $k \geq 3$  and for  $m \geq k$ , all  $k$ -homogeneous latin trades of volume  $km$  exist except possibly for finitely many  $m$ . We also show that for  $3 \leq k \leq 8$  and  $m \geq k$ ,  $k$ -homogeneous latin trades of volume  $km$  exist. It is obvious that we can omit empty rows and columns of a latin trade. Hence without loss of generality, we can assume any  $k$ -homogeneous latin trade of volume  $km$  is located in an  $m \times m$  square.

## 2 Results

For each  $k$ , there exists at least a  $k$ -homogeneous latin trade of volume  $k^2$ . To see this, for a latin square  $L$  of order  $k$ , we can take  $L^*$  to be a latin square with a cyclic permutation on the rows of  $L$ . So  $L^*$  is a disjoint mate of  $L$ .

**Theorem 1** *If  $l \neq 2, 6$  and for each  $k \in \{k_1, \dots, k_l\}$  there exists a  $k$ -homogeneous latin trade of volume  $kp$ , then a  $(k_1 + \dots + k_l)$ -homogeneous latin trade of volume  $(k_1 + \dots + k_l)lp$  exists. (Some  $k_i$ s can possibly be zero).*

**Proof.** Since  $l \neq 2, 6$ , there exist two  $l \times l$  orthogonal latin squares. Denote one of them by  $L$  and partition  $L$  into  $l$  transversals. Consider disjoint sets  $A_1, A_2, \dots, A_l$ , each with  $p$  elements. Then, if  $k_i \neq 0$  replace the element  $j$  in the  $i$ -th transversal of  $L$  by a  $k_i$ -homogeneous latin trade of volume  $k_i p$  with elements taken from  $A_j$ , and if  $k_i = 0$ , then we replace each element of  $i$ -th transversal by an empty  $p \times p$  square. ■

A latin square is *self-orthogonal* if it is orthogonal to its transpose. A self-orthogonal latin square (SOLS) of order  $n$  will be denoted by SOLS( $n$ ).

**Theorem 2** *For each  $k > 2$ , a  $k$ -homogeneous latin trade of volume  $k(k+1)$  exists.*

**Proof.** By Theorem 6.1.3 of [1], page 139, we know that SOLS( $n$ ) exists for every  $n \neq 2, 3$ , and 6. By deleting the main diagonals in an SOLS( $n$ ) and in its transpose, we obtain  $(n-1)$ -homogeneous latin trade of volume  $n(n-1)$  with its disjoint mate. An example of a 5-homogeneous latin trade of volume 30 is shown in Figure 2.

·	3 <sub>2</sub>	2 <sub>3</sub>	0 <sub>4</sub>	4 <sub>5</sub>	5 <sub>0</sub>
5 <sub>2</sub>	4 <sub>1</sub>	0 <sub>4</sub>	·	1 <sub>0</sub>	2 <sub>5</sub>
4 <sub>3</sub>	1 <sub>4</sub>	·	2 <sub>0</sub>	3 <sub>1</sub>	0 <sub>2</sub>
0 <sub>4</sub>	5 <sub>3</sub>	4 <sub>0</sub>	1 <sub>5</sub>	·	3 <sub>1</sub>
2 <sub>5</sub>	·	3 <sub>1</sub>	4 <sub>2</sub>	5 <sub>4</sub>	1 <sub>3</sub>
3 <sub>0</sub>	2 <sub>5</sub>	1 <sub>2</sub>	5 <sub>1</sub>	0 <sub>3</sub>	·

Figure 2: A 5-homogeneous latin trade of volume 30.


And this example completes the proof. ■

**Theorem 3** *Any 2-homogeneous latin trade  $T$  can be partitioned into disjoint intercalates.*

**Proof.** We prove by induction. Suppose  $T$  is a 2-homogeneous latin trade of volume  $2m$ . Without loss of generality, let  $\{(0, 0; 0), (0, 1; 1), (1, 0; 1)\} \subseteq T$ . Then  $T$  must contain  $(1, 1; 0)$ . We can apply the same argument to the  $(m-2) \times (m-2)$  subsquare obtained by removing rows 0 and 1, and columns 0 and 1. This completes the proof. ■

**Theorem 4** *For every  $k$ , if there exists a  $k$ -homogeneous latin trade of volume  $km$  and a  $k$ -homogeneous latin trade of volume  $kn$ , then for each  $r$  and  $s \geq 0$ , there exists a  $k$ -homogeneous latin trade of volume  $k(rm + sn)$ .*

**Proof.** Let  $T_1, \dots, T_r$  be  $k$ -homogeneous latin trades of volume  $km$  and  $T_{r+1}, \dots, T_{r+s}$  be  $k$ -homogeneous latin trades of volume  $kn$  such that for each  $i$ ,  $1 \leq i \leq r$  elements of  $T_i$  are in the set  $\{(i-1)m+1, \dots, im\}$  and for each  $j$ ,  $1 \leq j \leq s$  elements of  $T_{j+r}$  are in the set  $\{rm+(j-1)n+1, \dots, rm+jn\}$ . The following latin trade is a  $k$ -homogeneous latin trade of volume  $k(rm + sn)$ .



■

**Corollary 1** *For each  $k$  and  $m$  where  $m \geq k^2$ , there exists a  $k$ -homogeneous latin trade of volume  $km$ .*

**Proof.** If  $m \geq k^2$ , then we can write  $m$  as  $m = rk + s(k+1)$ , where  $r, s \geq 0$ . Theorem 4 and Theorem 2 lead us to conclusion. ■

**Theorem 5** *If  $(m, k) = d$ ,  $m \geq k$ , and  $d > 1$ , then there exists a  $k$ -homogeneous latin trade of volume  $km$ .*

**Proof.** For  $m = k$ , the theorem is trivial. Now suppose that  $m \neq k$ , so  $\frac{m}{d} \geq 2$ . Let  $m' = \frac{m}{d}$  and  $k' = \frac{k}{d}$ . We construct a  $k$ -homogeneous latin trade of volume  $km$  in the following way. Consider an  $m' \times m'$  latin square  $L$  on the set  $\{1, 2, \dots, m'\}$ . We replace each  $i$  in  $L$  with,

- a  $d$ -homogeneous latin trade of volume  $d^2$  whose elements are from the set  $\{(i-1)d+1, \dots, id\}$ , if  $1 \leq i \leq k'$ ; and
- an empty  $d \times d$  array, if  $k'+1 \leq i \leq m'$ .

So we obtain a  $k$ -homogeneous latin trade of volume  $km$ . Note that its mate can be obtained by replacing each  $d$ -homogeneous latin trade of volume  $d^2$  with its mate. In Figure 3, we have an example of the case  $k = 3d$  and  $m = 5d$ .

$T_1$	$T_2$	$T_3$	•	•
•	$T_1$	$T_2$	$T_3$	•
•	•	$T_1$	$T_2$	$T_3$
$T_3$	•	•	$T_1$	$T_2$
$T_2$	$T_3$	•	•	$T_1$

$T_1^*$	$T_2^*$	$T_3^*$	•	•
•	$T_1^*$	$T_2^*$	$T_3^*$	•
•	•	$T_1^*$	$T_2^*$	$T_3^*$
$T_3^*$	•	•	$T_1^*$	$T_2^*$
$T_2^*$	$T_3^*$	•	•	$T_1^*$

Figure 3: A latin trade is constructed for  $k' = 3$  and  $m' = 5$  ■

By using Theorem 5 and Theorem 3, we have the following corollary:

**Corollary 2** *For any  $m \geq 1$ , there exists a 2-homogeneous latin trade of volume  $2m$ , if and only if  $m$  is an even number.*

**Theorem 6** *For any  $m = 4l$  and  $2 \leq k \leq m$ , there exists a  $k$ -homogeneous latin trade of volume  $km$ .*

**Proof.** It is easy to see that theorem holds for  $l = 1$ . Also we may assume that  $m < k$ .

First, we prove the theorem in the case that  $l \neq 2, 6$ . We have the following cases to consider:

1.  $k$  is even. This case follows from Theorem 5.
2.  $k = 4l' + 1$ . This case follows from Theorem 1. Indeed, we set  $p = 4$  and  $k_i = 4$  for  $1 \leq i \leq l' - 1$ ,  $k_{l'} = 2$ ,  $k_{l'+1} = 3$ , and  $k_i = 0$  otherwise.
3.  $k = 4l' + 3$ . This case also follows from Theorem 1, by setting  $p = 4$  and  $k_i = 4$  for  $1 \leq i \leq l'$ ,  $k_{l'+1} = 3$ , and  $k_i = 0$  otherwise.

Next, we study the cases  $l = 2$  and  $l = 6$ . Actually for  $m = 8$  or  $24$ , the statement follows from Theorem 1, Theorem 2, Theorem 4 and Theorem 5, by choosing proper  $k_i$ s and  $p$ , except the case  $m = 8$  and  $k = 5$ . The existence of the last case is shown in Appendix. ■

**Theorem 7** *For any  $m = 5l$  and  $3 \leq k \leq m$ , there exists a  $k$ -homogeneous latin trade of volume  $km$ .*

**Proof.** Theorem trivially holds for  $l = 1$ . We may also assume that  $m > k$ .

First, we prove the theorem in the case that  $l \neq 2$  and  $6$ . We have the following cases to consider:

1.  $k = 5l'$ . Obviously this case follows from Theorem 5.
2.  $k = 5l' + 1$ . This case follows from Theorem 1. Indeed, we set  $k_i = 5$  for  $1 \leq i \leq l' - 1$  and  $k_{l'+1} = k_{l'} = 3$  and  $k_i = 0$  for  $l' + 2 \leq i \leq l$  and  $p = 5$ .
3.  $k = 5l' + 2$ . This case follows from Theorem 1, if we set  $k_i = 5$  for  $1 \leq i \leq l' - 1$  and  $k_{l'} = 3$  and  $k_{l'+1} = 4$  and  $k_i = 0$  for  $l' + 2 \leq i \leq l$  and  $p = 5$ .
4.  $k = 5l' + r$ ,  $r = 3, 4$ . This case also follows from Theorem 1, if we set  $k_i = 5$  for  $1 \leq i \leq l'$ ,  $k_{l'+1} = r$ , and  $k_i = 0$  for  $l' + 2 \leq i \leq l$ , and  $p = 5$ .

Next, for the case  $l = 2$ , by Theorem 1, Theorem 2, Theorem 4, and Theorem 5, we can show that there exists a  $k$ -homogeneous latin trade of volume  $10k$  for any  $3 \leq k \leq 10$  and  $k \neq 7$ . For  $k = 7$ , a 7-homogeneous latin trade of volume 70 is shown in Appendix.

If  $l = 6$ , by using Theorem 1, Theorem 2, Theorem 4 and Theorem 5, we can construct a  $k$ -homogeneous latin trade of volume  $30k$  for any  $3 \leq k \leq 30$ , by using  $k_i$ -homogeneous latin trades of volume  $6k_i$ , where  $k_1 + k_2 + k_3 + k_4 + k_5 = k$ ,  $0 \leq k_i \leq 6$  and  $k_i \neq 1$  for any  $i$ ,  $1 \leq i \leq 5$ . ■

**Theorem 8** *For any  $k \geq 3$  and  $m \geq 2k + 20$ , there exists a  $k$ -homogeneous latin trade of volume  $km$ .*

**Proof.** Consider an arbitrary  $m \geq 26$ . We can represent it as  $m = 4r + 5s$ , where  $r$  and  $s$  are positive integers. It is not hard to see that  $s \equiv m \pmod{4}$  and  $r \equiv 4m \pmod{5}$ . So we can conclude that, there exist unique  $0 \leq r' \leq 4$  and  $0 \leq s' \leq 3$ , such that  $r = 5a + r'$  and  $s = 4b + s'$ , where  $a, b \geq 0$ . It yields that  $m = 4r + 5s = 4(5a + r') + 5(4b + s')$ . We conclude that  $a + b = \frac{m - 4r' - 5s'}{20}$  is a constant number. Now we have two following cases:

- $a + b$  is even. In this case set  $a = b$ .
- $a + b = 2t + 1$ . If  $5s' > 4r'$ , set  $a = t + 1$  and  $b = t$ , otherwise set  $a = t$  and  $b = t + 1$ .

In each of these cases, we have  $|4r - 5s| \leq 20$ . And we have  $m = 4r + 5s$ , where  $4r, 5s \geq m/2 - 10$ . By Theorem 6 and Theorem 7, for any  $3 \leq k \leq m/2 - 10$  we have  $k$ -homogeneous latin trades of volume  $4kr$  and  $5ks$ . Now by Theorem 4, we conclude that there exists a  $k$ -homogeneous latin trade of volume  $4kr + 5ks$ . ■

The following theorem results immediately from Theorem 8.

**Main Theorem 1** For each given  $k \geq 3$ , and for  $m \geq k$ , all  $k$ -homogeneous latin trades of volume  $km$  exist except possibly for finitely many  $m$ .

**Theorem 9** Consider an arbitrary natural number  $k$ . If for any  $k + 1 \leq m \leq 2k - 1$  there exists a  $k$ -homogeneous latin trade of volume  $km$ , then for any  $m \geq k$  there exists a  $k$ -homogeneous latin trade of volume  $km$ .

**Proof.** For any  $m \geq 2k$ , we can write  $m = rk + sl$ , where  $r, s \geq 0$  and  $k + 1 \leq l \leq 2k - 1$ . Since there exist  $k$ -homogeneous latin trades of volume  $k^2$  and  $kl$ , by Theorem 4 we conclude that there exists a  $k$ -homogeneous latin trade of volume  $km$  ■.

**Main Theorem 2** For any  $3 \leq k \leq 8$  and  $m \geq k$ , there exists a  $k$ -homogeneous latin trade of volume  $km$ .

**Proof.** For  $k = 3$  or  $k = 4$  the theorem is proved in [4] and [3], respectively. By Theorem 9, we only need to show the existence of a  $k$ -homogeneous latin trade of volume  $km$ , for each  $5 \leq k \leq 8$  and  $k + 1 \leq m \leq 2k - 1$ . If  $m = k + 1$  statement follows from Theorem 2. For  $m > k + 1$ , we consider the following four cases:

- **Case 1.**  $k = 5$ .

$m = 7$ . An example of a 5-homogeneous latin trade of volume 35 is given in Appendix.

$m = 8$ . It follows from Theorem 6.

$m = 9$ . It follows from Theorem 1, by setting  $k_1 = 0$ ,  $k_2 = 2$ ,  $k_3 = 3$  and  $p = 3$ .

- **Case 2.**  $k = 6$ .

$m = 8, 9$  or  $10$ . All follow from Theorem 5.

$m = 11$ . An example of a 6-homogeneous latin trade of volume 66 is given in Appendix.

- **Case 3.**  $k = 7$ .

$m = 9$ . It follows from Theorem 1, by setting  $k_1 = 2$ ,  $k_2 = 2$ ,  $k_3 = 3$  and  $p = 3$ .

$m = 10$ . It follows from Theorem 7.

$m = 12$ . It follows from theorem 6.

$m = 11$  or  $m = 13$ . Examples of 7-homogeneous latin trades of volume 77 and 91 is given in Appendix.

- **Case 4.**  $k = 8$ .

$m = 10$ ,  $m = 12$  or  $m = 14$ . It follows from Theorem 5.

$m = 15$ . It follows from Theorem 7.

$m = 11$  or  $m = 13$ . There are examples of 8-homogeneous latin trades of volume 88 and 104: ■

### 3 Future Research

For future research, we could look at ensuring the latin trades we create are minimal. When we wish to determine if a set of entries is a subset of exactly one latin square, it is faster to use only minimal latin trades. Thus, we can look at the problem of whether minimal  $k$ -homogeneous latin trades of volume  $km$  exist for all  $k$  and  $m$ .

### References

- [1] Ian Anderson. *Combinatorial designs: Construction Methods*. John Wiley & Sons, Inc., New York, 1990.
- [2] Elizabeth J. Billington. Combinatorial trades: a survey of recent results. In *Designs, 2002*, volume 563 of *Math. Appl.*, pages 47–67. Kluwer Acad. Publ., Boston, MA, 2003.
- [3] Nicholas Cavenagh, Diane Donovan, and Aleš Drápal. 4-homogeneous latin trades. *Australasian Journal of Combinatorics*, to appear.
- [4] Nicholas J. Cavenagh, Aleš Drápal, and Diane Donovan. 3-homogeneous latin trades. *Preprint*.
- [5] A. D. Keedwell. Critical sets for Latin squares, graphs and block designs: a survey. *Congr. Numer.*, 113:231–245, 1996. Festschrift for C. St. J. A. Nash-Williams.
- [6] A. D. Keedwell. Critical sets in latin squares and related matters: an update. *Util. Math.*, 65:97–131, 2004.

## 4 Appendix

In the following all required  $k$ -homogeneous latin trades for the mentioned results are given.

5-homogeneous latin trades of volume 35 and 40:

2 <sub>0</sub>	4 <sub>2</sub>	0 <sub>1</sub>	3 <sub>4</sub>	1 <sub>3</sub>	·	·
3 <sub>2</sub>	5 <sub>1</sub>	2 <sub>0</sub>	0 <sub>5</sub>	·	1 <sub>3</sub>	·
·	·	4 <sub>2</sub>	5 <sub>6</sub>	2 <sub>5</sub>	3 <sub>4</sub>	6 <sub>3</sub>
·	2 <sub>5</sub>	1 <sub>6</sub>	6 <sub>3</sub>	·	5 <sub>1</sub>	3 <sub>2</sub>
6 <sub>3</sub>	1 <sub>6</sub>	·	4 <sub>0</sub>	3 <sub>4</sub>	·	0 <sub>1</sub>
5 <sub>6</sub>	·	6 <sub>4</sub>	·	4 <sub>2</sub>	0 <sub>5</sub>	2 <sub>0</sub>
0 <sub>5</sub>	6 <sub>4</sub>	·	·	5 <sub>1</sub>	4 <sub>0</sub>	1 <sub>6</sub>

7 <sub>0</sub>	0 <sub>1</sub>	3 <sub>2</sub>	2 <sub>3</sub>	·	·	·	1 <sub>7</sub>
0 <sub>1</sub>	1 <sub>0</sub>	4 <sub>3</sub>	3 <sub>2</sub>	·	2 <sub>4</sub>	·	·
·	·	·	7 <sub>1</sub>	1 <sub>6</sub>	6 <sub>7</sub>	5 <sub>4</sub>	4 <sub>5</sub>
·	4 <sub>2</sub>	·	·	6 <sub>7</sub>	7 <sub>6</sub>	2 <sub>5</sub>	5 <sub>4</sub>
·	2 <sub>5</sub>	5 <sub>6</sub>	6 <sub>7</sub>	7 <sub>0</sub>	·	0 <sub>2</sub>	·
6 <sub>5</sub>	5 <sub>4</sub>	·	1 <sub>6</sub>	3 <sub>1</sub>	·	4 <sub>3</sub>	·
1 <sub>6</sub>	·	6 <sub>4</sub>	·	·	4 <sub>3</sub>	3 <sub>0</sub>	0 <sub>1</sub>
5 <sub>7</sub>	·	2 <sub>5</sub>	·	0 <sub>3</sub>	3 <sub>2</sub>	·	7 <sub>0</sub>

A 6-homogeneous latin trade of volume 66 and a 7-homogeneous latin trade of volume 70:

·	·	6 <sub>2</sub>	·	·	7 <sub>5</sub>	5 <sub>6</sub>	a <sub>7</sub>	2 <sub>8</sub>	·	8 <sub>a</sub>
9 <sub>1</sub>	·	·	7 <sub>4</sub>	·	·	a <sub>7</sub>	1 <sub>8</sub>	8 <sub>9</sub>	4 <sub>a</sub>	·
·	6 <sub>3</sub>	·	0 <sub>5</sub>	3 <sub>6</sub>	8 <sub>7</sub>	7 <sub>8</sub>	·	·	5 <sub>0</sub>	·
8 <sub>3</sub>	·	2 <sub>5</sub>	·	·	1 <sub>8</sub>	·	5 <sub>a</sub>	·	a <sub>1</sub>	3 <sub>2</sub>
1 <sub>4</sub>	·	7 <sub>6</sub>	4 <sub>7</sub>	·	·	6 <sub>a</sub>	·	3 <sub>1</sub>	·	a <sub>3</sub>
·	7 <sub>6</sub>	9 <sub>7</sub>	·	1 <sub>9</sub>	·	·	2 <sub>1</sub>	4 <sub>2</sub>	·	6 <sub>4</sub>
·	3 <sub>7</sub>	·	2 <sub>9</sub>	·	4 <sub>0</sub>	·	7 <sub>2</sub>	9 <sub>3</sub>	0 <sub>4</sub>	·
·	·	0 <sub>9</sub>	·	9 <sub>0</sub>	5 <sub>1</sub>	·	·	1 <sub>4</sub>	6 <sub>5</sub>	4 <sub>6</sub>
3 <sub>8</sub>	0 <sub>9</sub>	·	9 <sub>0</sub>	6 <sub>1</sub>	·	8 <sub>3</sub>	·	·	1 <sub>6</sub>	·
a <sub>9</sub>	9 <sub>a</sub>	5 <sub>0</sub>	·	0 <sub>2</sub>	·	·	8 <sub>5</sub>	·	·	2 <sub>8</sub>
4 <sub>a</sub>	a <sub>0</sub>	·	5 <sub>2</sub>	2 <sub>3</sub>	0 <sub>4</sub>	3 <sub>5</sub>	·	·	·	·

·	·	·	3 <sub>2</sub>	1 <sub>3</sub>	5 <sub>9</sub>	2 <sub>5</sub>	7 <sub>1</sub>	8 <sub>7</sub>	9 <sub>8</sub>
·	·	·	7 <sub>3</sub>	5 <sub>4</sub>	3 <sub>5</sub>	4 <sub>6</sub>	8 <sub>7</sub>	9 <sub>8</sub>	6 <sub>9</sub>
·	5 <sub>2</sub>	·	·	7 <sub>5</sub>	0 <sub>1</sub>	1 <sub>7</sub>	9 <sub>8</sub>	2 <sub>9</sub>	8 <sub>0</sub>
1 <sub>2</sub>	9 <sub>3</sub>	5 <sub>4</sub>	0 <sub>5</sub>	·	·	·	3 <sub>9</sub>	4 <sub>0</sub>	2 <sub>1</sub>
2 <sub>3</sub>	3 <sub>4</sub>	7 <sub>5</sub>	5 <sub>6</sub>	0 <sub>7</sub>	·	·	6 <sub>0</sub>	·	4 <sub>2</sub>
5 <sub>9</sub>	4 <sub>5</sub>	9 <sub>1</sub>	2 <sub>7</sub>	·	6 <sub>4</sub>	·	1 <sub>6</sub>	7 <sub>2</sub>	·
7 <sub>5</sub>	8 <sub>6</sub>	0 <sub>7</sub>	6 <sub>8</sub>	·	4 <sub>0</sub>	5 <sub>1</sub>	·	·	1 <sub>4</sub>
8 <sub>1</sub>	·	4 <sub>8</sub>	·	2 <sub>0</sub>	1 <sub>6</sub>	3 <sub>2</sub>	0 <sub>3</sub>	6 <sub>4</sub>	·
9 <sub>7</sub>	6 <sub>8</sub>	1 <sub>9</sub>	8 <sub>0</sub>	3 <sub>1</sub>	·	7 <sub>3</sub>	·	·	0 <sub>6</sub>
3 <sub>8</sub>	2 <sub>9</sub>	8 <sub>0</sub>	·	4 <sub>2</sub>	9 <sub>3</sub>	6 <sub>4</sub>	·	0 <sub>6</sub>	·

7-homogeneous latin trades of volume 77 and 91

·	·	·	·	6 <sub>4</sub>	4 <sub>5</sub>	9 <sub>6</sub>	8 <sub>7</sub>	a <sub>8</sub>	5 <sub>9</sub>	7 <sub>a</sub>
·	·	·	8 <sub>4</sub>	·	7 <sub>6</sub>	6 <sub>7</sub>	a <sub>8</sub>	0 <sub>9</sub>	9 <sub>a</sub>	4 <sub>0</sub>
·	·	·	7 <sub>5</sub>	0 <sub>6</sub>	6 <sub>7</sub>	5 <sub>8</sub>	·	8 <sub>a</sub>	1 <sub>0</sub>	a <sub>1</sub>
5 <sub>3</sub>	9 <sub>4</sub>	0 <sub>5</sub>	·	·	·	3 <sub>9</sub>	1 <sub>a</sub>	4 <sub>0</sub>	a <sub>1</sub>	·
6 <sub>4</sub>	7 <sub>5</sub>	5 <sub>6</sub>	4 <sub>7</sub>	·	·	·	2 <sub>0</sub>	·	3 <sub>2</sub>	0 <sub>3</sub>
4 <sub>5</sub>	5 <sub>6</sub>	·	1 <sub>8</sub>	·	·	·	6 <sub>1</sub>	3 <sub>2</sub>	2 <sub>3</sub>	8 <sub>4</sub>
8 <sub>6</sub>	6 <sub>7</sub>	a <sub>8</sub>	2 <sub>9</sub>	3 <sub>a</sub>	·	·	7 <sub>2</sub>	9 <sub>3</sub>	·	·
9 <sub>7</sub>	·	1 <sub>9</sub>	·	4 <sub>0</sub>	5 <sub>1</sub>	7 <sub>2</sub>	·	2 <sub>4</sub>	0 <sub>5</sub>	·
7 <sub>8</sub>	a <sub>9</sub>	9 <sub>a</sub>	·	2 <sub>1</sub>	1 <sub>2</sub>	8 <sub>3</sub>	·	·	·	3 <sub>7</sub>
3 <sub>9</sub>	0 <sub>a</sub>	8 <sub>0</sub>	9 <sub>1</sub>	a <sub>2</sub>	2 <sub>3</sub>	·	·	·	·	1 <sub>8</sub>
·	4 <sub>0</sub>	6 <sub>1</sub>	5 <sub>2</sub>	1 <sub>3</sub>	3 <sub>4</sub>	2 <sub>5</sub>	0 <sub>6</sub>	·	·	·

·	·	·	·	9 <sub>4</sub>	·	7 <sub>6</sub>	4 <sub>7</sub>	a <sub>8</sub>	c <sub>9</sub>	6 <sub>a</sub>	·	8 <sub>c</sub>
·	·	·	9 <sub>4</sub>	·	0 <sub>6</sub>	6 <sub>7</sub>	·	7 <sub>9</sub>	b <sub>a</sub>	a <sub>b</sub>	·	4 <sub>0</sub>
·	·	·	·	a <sub>6</sub>	·	·	6 <sub>9</sub>	b <sub>a</sub>	9 <sub>b</sub>	0 <sub>c</sub>	1 <sub>0</sub>	c <sub>1</sub>
5 <sub>3</sub>	·	c <sub>5</sub>	·	·	·	·	·	2 <sub>b</sub>	3 <sub>c</sub>	b <sub>0</sub>	0 <sub>1</sub>	1 <sub>2</sub>
·	6 <sub>5</sub>	1 <sub>6</sub>	b <sub>7</sub>	·	·	·	7 <sub>b</sub>	·	·	5 <sub>1</sub>	3 <sub>2</sub>	2 <sub>3</sub>
9 <sub>5</sub>	0 <sub>6</sub>	·	4 <sub>8</sub>	6 <sub>9</sub>	·	·	·	8 <sub>0</sub>	·	·	5 <sub>3</sub>	3 <sub>4</sub>
·	b <sub>7</sub>	9 <sub>8</sub>	a <sub>9</sub>	4 <sub>a</sub>	2 <sub>b</sub>	·	·	·	7 <sub>2</sub>	·	8 <sub>4</sub>	·
3 <sub>7</sub>	7 <sub>8</sub>	5 <sub>9</sub>	8 <sub>a</sub>	·	·	·	·	9 <sub>2</sub>	a <sub>3</sub>	·	2 <sub>5</sub>	·
a <sub>8</sub>	·	8 <sub>a</sub>	1 <sub>b</sub>	0 <sub>c</sub>	b <sub>0</sub>	5 <sub>1</sub>	·	·	·	c <sub>5</sub>	·	·
8 <sub>9</sub>	·	·	·	2 <sub>0</sub>	6 <sub>1</sub>	3 <sub>2</sub>	9 <sub>3</sub>	·	·	1 <sub>6</sub>	·	0 <sub>8</sub>
c <sub>a</sub>	8 <sub>b</sub>	a <sub>c</sub>	·	·	3 <sub>2</sub>	2 <sub>3</sub>	b <sub>4</sub>	·	·	·	·	4 <sub>8</sub>
·	5 <sub>c</sub>	·	7 <sub>1</sub>	c <sub>2</sub>	4 <sub>3</sub>	1 <sub>4</sub>	3 <sub>5</sub>	·	2 <sub>7</sub>	·	·	·
7 <sub>c</sub>	c <sub>0</sub>	6 <sub>1</sub>	·	·	1 <sub>4</sub>	4 <sub>5</sub>	5 <sub>6</sub>	0 <sub>7</sub>	·	·	·	·

8-homogeneous latin trades of volume 88 and 104:

·	·	·	4 <sub>3</sub>	3 <sub>4</sub>	6 <sub>5</sub>	5 <sub>6</sub>	8 <sub>7</sub>	9 <sub>8</sub>	a <sub>9</sub>	7 <sub>a</sub>
·	·	·	5 <sub>4</sub>	0 <sub>5</sub>	4 <sub>6</sub>	6 <sub>7</sub>	7 <sub>8</sub>	8 <sub>9</sub>	9 <sub>a</sub>	a <sub>0</sub>
·	·	·	7 <sub>5</sub>	1 <sub>6</sub>	5 <sub>7</sub>	9 <sub>8</sub>	a <sub>9</sub>	0 <sub>a</sub>	6 <sub>0</sub>	8 <sub>1</sub>
4 <sub>3</sub>	0 <sub>4</sub>	a <sub>5</sub>	·	·	·	2 <sub>9</sub>	9 <sub>a</sub>	1 <sub>0</sub>	5 <sub>1</sub>	3 <sub>2</sub>
5 <sub>4</sub>	6 <sub>5</sub>	7 <sub>6</sub>	2 <sub>7</sub>	·	·	·	1 <sub>0</sub>	3 <sub>1</sub>	0 <sub>2</sub>	4 <sub>3</sub>
7 <sub>5</sub>	5 <sub>6</sub>	8 <sub>7</sub>	3 <sub>8</sub>	·	·	·	6 <sub>1</sub>	4 <sub>2</sub>	2 <sub>3</sub>	1 <sub>4</sub>
3 <sub>6</sub>	9 <sub>7</sub>	6 <sub>8</sub>	8 <sub>9</sub>	2 <sub>a</sub>	7 <sub>0</sub>	·	0 <sub>2</sub>	a <sub>3</sub>	·	·
9 <sub>7</sub>	7 <sub>8</sub>	5 <sub>9</sub>	·	4 <sub>0</sub>	0 <sub>1</sub>	8 <sub>2</sub>	·	2 <sub>4</sub>	1 <sub>5</sub>	·
a <sub>8</sub>	8 <sub>9</sub>	9 <sub>a</sub>	·	6 <sub>1</sub>	1 <sub>2</sub>	7 <sub>3</sub>	·	·	3 <sub>6</sub>	2 <sub>7</sub>
8 <sub>9</sub>	4 <sub>a</sub>	1 <sub>0</sub>	9 <sub>1</sub>	a <sub>2</sub>	2 <sub>3</sub>	3 <sub>4</sub>	·	·	·	0 <sub>8</sub>
6 <sub>a</sub>	a <sub>0</sub>	0 <sub>1</sub>	1 <sub>2</sub>	5 <sub>3</sub>	3 <sub>4</sub>	4 <sub>5</sub>	2 <sub>6</sub>	·	·	·

·	·	·	·	·	b <sub>5</sub>	5 <sub>6</sub>	8 <sub>7</sub>	7 <sub>8</sub>	a <sub>9</sub>	6 <sub>a</sub>	c <sub>b</sub>	9 <sub>c</sub>
·	·	·	b <sub>4</sub>	·	·	8 <sub>7</sub>	9 <sub>8</sub>	4 <sub>9</sub>	7 <sub>a</sub>	a <sub>b</sub>	0 <sub>c</sub>	c <sub>0</sub>
·	·	·	·	a <sub>6</sub>	·	9 <sub>8</sub>	6 <sub>9</sub>	b <sub>a</sub>	c <sub>b</sub>	0 <sub>c</sub>	1 <sub>0</sub>	8 <sub>1</sub>
c <sub>3</sub>	·	b <sub>5</sub>	·	·	·	3 <sub>9</sub>	·	2 <sub>b</sub>	9 <sub>c</sub>	1 <sub>0</sub>	5 <sub>1</sub>	0 <sub>2</sub>
5 <sub>4</sub>	7 <sub>5</sub>	0 <sub>6</sub>	1 <sub>7</sub>	·	·	·	·	·	6 <sub>0</sub>	2 <sub>1</sub>	3 <sub>2</sub>	4 <sub>3</sub>
8 <sub>5</sub>	9 <sub>6</sub>	6 <sub>7</sub>	7 <sub>8</sub>	3 <sub>9</sub>	·	·	·	·	·	5 <sub>2</sub>	4 <sub>3</sub>	2 <sub>4</sub>
4 <sub>6</sub>	8 <sub>7</sub>	7 <sub>8</sub>	a <sub>9</sub>	6 <sub>a</sub>	1 <sub>b</sub>	·	·	9 <sub>1</sub>	·	·	b <sub>4</sub>	·
3 <sub>7</sub>	c <sub>8</sub>	·	8 <sub>a</sub>	·	5 <sub>c</sub>	7 <sub>0</sub>	·	a <sub>2</sub>	0 <sub>3</sub>	·	2 <sub>5</sub>	·
a <sub>8</sub>	b <sub>9</sub>	8 <sub>a</sub>	9 <sub>b</sub>	0 <sub>c</sub>	2 <sub>0</sub>	·	5 <sub>2</sub>	·	·	c <sub>5</sub>	·	·
·	6 <sub>a</sub>	a <sub>b</sub>	·	1 <sub>0</sub>	0 <sub>1</sub>	·	4 <sub>3</sub>	8 <sub>4</sub>	·	b <sub>6</sub>	·	3 <sub>8</sub>
6 <sub>a</sub>	a <sub>b</sub>	·	·	9 <sub>1</sub>	3 <sub>2</sub>	4 <sub>3</sub>	2 <sub>4</sub>	·	b <sub>6</sub>	·	·	1 <sub>9</sub>
·	5 <sub>c</sub>	1 <sub>0</sub>	2 <sub>1</sub>	c <sub>2</sub>	4 <sub>3</sub>	0 <sub>4</sub>	7 <sub>5</sub>	·	3 <sub>7</sub>	·	·	·
7 <sub>c</sub>	·	5 <sub>1</sub>	4 <sub>2</sub>	2 <sub>3</sub>	c <sub>4</sub>	6 <sub>5</sub>	3 <sub>6</sub>	1 <sub>7</sub>	·	·	·	·